

CLAIM AMENDMENTS:

1. (previously ~~amended~~ presented) An aqueous hydrocarbon fuel emulsion comprising water, fuel, and an emulsifier comprising an amino alkylphenol which is made by reacting alkylphenol, an aldehyde and an amine in a ratio of 1:0:9:0.1 to 1:1:9:1.9 molar resulting in an amino alkyl phenol.

2 (original) The emulsion of claim 1 wherein the amino alkylphenol is in combination with an emulsifier selected from the group consisting of at least one of:

(i) at least one fuel-soluble product made by reacting at least one hydrocarbyl-substituted carboxylic acid acylating agent with ammonia or an amine, the hydrocarbyl substituent of said acylating agent having about 50 to about 500 carbon atoms;

(ii) at least one of an ionic or a nonionic compound having a hydrophilic-lipophilic balance (HLB) of about 1 to about 40;

(iii) mixture of (ii) with (i); and

(iv) a water-soluble compound selected from the group consisting of amine salts, ammonium salts, azide compounds, nitrate esters, nitramine, nitro compounds, alkali metal salts, alkaline earth metal salts, in combination with (i), (ii), (iii) or (v);

(v) the reaction product of polyacidic polymer with at least one fuel soluble product made by reacting at least one hydrocarbyl-substituted carboxylic acid acylating agent with ammonia, an amine or a polyamine;

and combinations thereof.

3. (original) The emulsion of claim 1 wherein the amino alkylphenol is made by the reaction selected from the group consisting of (a) the reaction of an alkylphenol directly with an aldehyde and an amine resulting in an alkylphenol monomer connected by a methylene group to an amine; (b) the reaction of an alkylphenol with an aldehyde resulting in an oligomer wherein the alkylphenols are bridged with methylene groups, the oligomer is then reacted with more aldehyde and an amine to give a Mannich product; and (c) combinations of (a) and (b).

4. (original) The emulsion of claim 1 wherein the alkylphenols have an alkyl group selected from C₆ to C₁₇₀ and wherein the alkyl group is linear, branched or a combination thereof.

5. (original) The emulsion of claim 1 wherein the alkylphenols are selected from the group consisting of polypropylphenol, polybutylphenol, poly(isobutenyl)phenol, polyamylphenol, tetrapropylphenol, substituted phenols, and combinations thereof.


6. (original) The emulsion of claim 5 wherein the amino alkylphenol is selected from the group consisting of tetrapropenylphenol, poly(isobutenyl)phenol, and combinations thereof.

7. (original) The emulsion of claim 1 wherein the aldehydes are selected from the group consisting of aliphatic aldehydes, including, but not limited to, formaldehyde; acetaldehyde; aldol (β -hydroxy butyraldehyde); aromatic aldehydes, selected from the group consisting of benzaldehyde; heterocyclic aldehydes, selected from the group consisting of furfural, and combinations thereof.


8. (original) The emulsion of claim 7 wherein the aldehyde is formaldehyde.

9. (previously ~~amended~~ presented) The emulsion of claim 1 wherein the amine is selected from the group consisting of alkanolamines selected from the group consisting of monoethanol amine, diethanolamine, N-(2-aminoethyl) ethanolamine, and combinations thereof; di- and polyamines selected from the group consisting of polyalkylene amines, dimethylaminopropylamine, 3-aminopropyl morpholine, ethylenediamine, diethylenetriamine, triethylene tetramine, tetraethylene pentamine, and combinations thereof; distillation bottoms; polyalkyl polyamines; propylene diamine; aromatic amines selected from the group consisting of o-, m- and p-phenylene diamine, diamino naphthalenes; acid-substituted polyalkylpolyamines selected from the group consisting of N-acetyl tetraethylenepentamine, and the corresponding

formyl-, propionyl-, butyl-, N-substituted compounds, and combinations thereof; cyclized N-compounds selected from the group consisting N-alkyl amines of imidazolidine, pyrimidine, and combinations thereof; morpholine, thiomorpholine, pyrrole, pyrroline, pyrrolidine, indole, pyrazole, pyrazoline, pyrazolidine, imidazole, imidazoline, imidazolidine, piperidine, phenoxazine, phenthiazine and their substituted analogs, the product obtained by reacting an alkenyl succinic anhydride of the formula



or alkenyl succinic acid of the formula



and combinations thereof

wherein R is an alkylene group.

10. (original) The emulsion of claim 1 wherein the emulsion comprises the emulsifier in the range of about 0.05% to about 20% by weight of the water fuel emulsion, the fuel in the range of about 50% to about 95 % by weight of the water fuel emulsion, and the water in the range of about 1% to about 50% by weight of the water fuel emulsion, and wherein the emulsion has a mean particle droplet in the range of about 0.1 micron to about 1 micron.

11. (currently amended) A process for making an aqueous hydrocarbon fuel comprising

(a) mixing a liquid hydrocarbon fuel and at least one emulsifier to form a hydrocarbon fuel emulsifier mixture wherein at least one emulsifier comprises an amino alkylphenol which is the reaction product of an alkylphenol, an aldehyde and an amine in a ratio range of 1:0.9:0.1 to 1:1.9:1.9 molar; and

(b) mixing the hydrocarbon fuel emulsifier mixture with water or water and ammonium nitrate under emulsification conditions to form an aqueous hydrocarbon fuel composition, wherein the aqueous hydrocarbon fuel composition includes a discontinuous phase, the discontinuous aqueous phase being comprised of aqueous droplets having a mean diameter of 1.0 micron or less.

12. (original) The emulsion of claim 11 wherein the amino alkylphenol is made by the reaction selected from the group consisting of (a) the reaction of an alkylphenol directly with an aldehyde and an amine resulting in an alkylphenol monomer connected by a methylene group to an amine; (b) the reaction of an alkylphenol with an aldehyde resulting in an oligomer wherein the alkylphenols are bridged with methylene groups, the oligomer is then reacted with more aldehyde and an amine to give a Mannich product; and (c) combinations of (a) and (b).

13. (original) The process of claim 12 wherein the amino alkylphenol emulsifier has been made by reacting the alkylphenol:aldehyde:amine in reaction (a) in a ratio range of 1:1:0.1 molar to 1:2:2 molar.

14. (original) The process of claim 13 wherein the amino alkylphenol emulsifier has been made by reacting the alkylphenol:aldehyde:amine in reaction (a) in a ratio range of 1:0.9:0.1 to 1:1.9:1.9 molar.

15. (original) The process of claim 14 wherein the amino alkylphenol emulsifier has been made by reacting the alkylphenol:aldehyde:amine in reaction (a) in a ratio range of 1:1.5:1.2 to 1:1.9:1.8 molar.

16. (original) The process of claim 15 wherein the amino alkylphenol emulsifier has been made by reacting the alkylphenol:aldehyde:amine in reaction (a) in a ratio range of 1:0.8:0.3 to 1:1.5:0.7 molar.

17. (original) The process of claim 11 wherein the reaction is carried out under conditions to provide the formation of the desired product wherein the reaction temperature is in the range of 40°C to about 200°C and the pressure in the range of elevated pressure to reduce pressure and it occurs over a period of time in the range of about 15 minutes to about 8 hours.

18. (original) The process of claim 11 wherein the concentration of the emulsifier in the water-blend fuel is in the range of about 0.05% to about 20% by weight of the total emulsion.

19. (original) The process of claim 12 wherein the reaction (b) is carried out at a temperature in the range of about 0°C to about 150°C for a period of time ranging from 15 minutes to about 8 hours resulting in the oligomer wherein the alkylphenols are bridged with a methylene group; the intermediate product is reacted in the range of about 1 mole oligomer:about 0.1 mole amine to about 1 mole oligomer:about 2 moles amine.

20. (original) The process of claim 19 wherein the intermediate product is reacted in the range of about 1 mole oligomer:about 0.2 mole amine to about 1 mole oligomer:about 1.5 moles amine.

21. (original) The process of claim 20 wherein the intermediate product is reacted in the range of about 1 mole oligomer:about 0.3 mole amine to about 1 mole oligomer:about 0.9 moles amine.

22. (original) A method for fueling an engine comprising fueling the engine with the composition of claim 1.

RESPONSE AND REQUEST FOR RECONSIDERATION

Please enter the amendment as shown. Applicants have amended claims 1 and 9 to correctly state the status identifier. Thus, claims 1-22 are pending in the application. The current status of all claims is shown in the claim amendment section of this amendment.


The amendment to claims 1 and 9 obviates the notice of non-compliance. Accordingly, Applicants submission of the entire Amendments to the claims is in compliance with 37 CFR 1.121.

For the foregoing reasons, it is submitted that the present claims are in condition for allowance. The foregoing remarks are believed to be a full and complete response to the outstanding office action. Therefore, favorable reconsideration is respectfully requested. If for any reason the Examiner believes a telephone conference would expedite the prosecution of this application, it is respectfully requested that she call Applicants' attorney at (440) 347-5072.

Any required fees or any deficiency or overpayment in fees should be charged or credited to the Lubrizol Corporation Deposit Account No. 12-2275.

Respectfully submitted,

THE LUBRIZOL CORPORATION



Teresan W. Gilbert
Attorney for Applicant
Reg. No. 31,360

The Lubrizol Corporation
29400 Lakeland Blvd.
Wickliffe, Ohio 44092-2298
Telephone: (440) 347-5072
Facsimile: (440) 347-1110